arXiv:cond-mat/0502251v1 [cond-mat.other] 9 Feb 2005

# Cold atoms in non-Abelian gauge potentials: From the Hofstadter "moth" to lattice gauge theory

K. Osterloh[1], M. Baig[2], L. Santos[3], P. Zoller[4], and M. Lewenstein[1,5]
*(1) Institut für Theoretische Physik, Universität Hannover, D-30167 Hannover, Germany*
*(2) IFAE, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain*
*(3) Institut für Theoretische Physik III, Universität Stuttgart, D-70550 Stuttgart*
*(4) Institute for Theoretical Physics, University of Innsbruck, A-6020 Innsbruck, Austria*
*(5) ICFO — Institut de Ciències Fotòniques, 08034 Barcelona, Spain*
(Dated: February 2, 2008)

We demonstrate how to create artificial external non-Abelian gauge potentials acting on cold atoms in optical lattices. The method employs $n$ internal states of atoms and laser assisted state sensitive tunneling. Thus, dynamics are communicated by unitary $n\times n$-matrices. By experimental control of the tunneling parameters, the system can be made truly non-Abelian. We show that single particle dynamics in the case of intense $U(2)$ vector potentials lead to a generalized Hofstadter butterfly spectrum which shows a complex "moth"-like structure. We discuss the possibility to employ non-Abelian interferometry (Aharonov-Bohm effect) and address methods to realize matter dynamics in specific classes of lattice gauge fields.

One of the most significant trends in the physics of ultra-cold gases nowadays concerns, without any doubts, strongly correlated systems. Apart from systems at BCS-BEC crossover, low dimensional systems, and atomic lattice gases, perhaps the most fascinating possibilities are offered by atomic gases subject to the effects of artificial magnetic fields. A feasible realization, albeit experimentally challenging, is an atomic system in a rotating trap. Trap rotation mimics the effects of external homogeneous magnetic fields. In the situation, when the rotation frequency approaches the trap frequency, the systems may allow to study the fractional quantum Hall effect (FQHE) and Laughlin liquids [1]. Several experiments with gases in rotating traps are currently underway [2]. In this context, rotating dipolar lattice gases (for a review, see [3]) offer new possibilities and advantages compared to gases with short ranged interactions, as the energy gap between the Laughlin state and quasi-hole excited states survives the large $N$ limit. This allows for the realization of Laughlin liquids with a few hundred atoms at realistic temperatures [4]. Very recently, the creation of artificial magnetic fields in terms of electro-magnetically induced transparency (EIT) has been proposed in Ref. [8].

Alternatively, magnetic field effects can be realized in a lattice by introducing appropriate phase factors for tunneling amplitudes. If multiplied around an elementary plaquette, these lead to phases proportional to the magnetic flux penetrating the plaquette. Jaksch and Zoller, and subsequently other groups, have recently proposed methods to realize such "artificial" magnetic field effects in lattice gases. These employ atoms with multiple internal states, laser assisted tunneling, lattice tilting (acceleration), and other experimentally accessible techniques [5, 6, 7]. The physical features in such "artificial" magnetic fields are extremely rich. For single atoms, the spectrum exhibits the fractal Hofstadter "butterfly" structure[9] illustrated in Fig. 1. In weakly interacting or weakly disordered systems, the modifications of the butterfly due to interactions and disorder, respectively, may be studied [5]. Finally, considering strong magnetic fields in the limit of strong interactions, Laughlin like states are expected [7]. In this Letter, we show that the use of atoms with more internal states, the application of state dependent, laser assisted tunelling (cf. [10]), and coherent transfer between internal states, allows for a generalization of the above methods and creation of "artificial external magnetic fields" corresponding to non-Abelian $U(n)$, or $SU(n)$ gauge fields [11]. In this case, tunneling amplitudes are replaced by unitary matrices whose product around a plaquette is non-trivial and its mean trace (Wilson loop) is not equal to $n$ [12]. To our knowledge, the physics of matter in designed non-Abelian gauge fields has not been studied at all in the AMO physics context. Although single particles in $SU(n)$ monopole fields have been intensively studied in high energy physics [13], other field configurations have not attracted such interest. Specific examples of external gauge fields have also been discussed with respect to NMR, nuclear and molecular physics [14]. Recently, there has been a proposal to create non-Abelian gauge fields using EIT [15]. However, all of these proposals either do not deal with interacting many particle systems in such gauge fields, or they do

FIG. 1: The Hofstadter butterfly. Eigenenergies in natural units plotted versus magnetic flux per plaquette $\alpha = p/q, \in [0, 1]$, with $q \leq 100$. The lowest Bloch band splits into subbands and forms a fractal for the set of all $\alpha$'s.



not consider lattice gauge fields of ultra-high strength.

Having introduced the designed external non-Abelian gauge fields in optical lattices, we study their influence on single particle dynamics. We generalize the Hofstadter butterfly to the non-Abelian case and show that in case of intense $U(2)$ gauge fields, the spectrum (as a function of two "magnetic fluxes") exhibits a complex "moth" structure of holes. We discuss the possibility to observe the $U(2)$ Aharonov-Bohm effect, to realize non-Abelian interferometry, and to simulate lattice gauge dynamics using random field configurations.

We start from the physics of a single particle in a two-dimensional square lattice of spacing $a$ in the presence of an Abelian magnetic field $B$. When the lattice potential is sufficiently strong, the single particle non-relativistic Hamiltonian in the tight binding approximation is given by the so-called Peierl's substitution [9], and reads

$$H=-2V_0\Big[\cos\Big(\frac{a}{\hbar}(p_x-\mathrm{i}\frac{\mathrm{e}}{c}A_x)\Big)+\cos\Big(\frac{a}{\hbar}(p_y-\mathrm{i}\frac{\mathrm{e}}{c}A_y)\Big)\Big], \quad (1)$$

where $V_0$ is the optical potential strength, $\vec{p}$ is the momentum operator, and $\vec{A}$ is the magnetic vector potential. Given $[p_j, A_j] = 0$, the single-particle wave equation reads

$$\begin{aligned} &e^{-\mathrm{i}\frac{\mathrm{e}a}{\hbar c}A_x}\psi(x+a,y)+e^{\mathrm{i}\frac{\mathrm{e}a}{\hbar c}A_x}\psi(x-a,y) \quad (2)\\ &+e^{-\mathrm{i}\frac{\mathrm{e}a}{\hbar c}A_y}\psi(x,y+a)+e^{\mathrm{i}\frac{\mathrm{e}a}{\hbar c}A_y}\psi(x,y-a)=-\frac{E}{V_0}\psi(x,\,y)\end{aligned}$$

The choice of $\vec{A}$ determines the magnetic field $\vec{B}$, which in turn determines the behavior of the system. For $\vec{B}=B\vec{\mathrm{e}}_z$, one may choose the potential $\vec{A}=(0,\,Bx,\,0)$, thus, solely tunnelings in the $y$-direction acquire phases. This makes the problem effectively one-dimensional and (2) can be transformed into Harper's equation [17]:

$$g(m+1)-g(m-1)+2\cos(2\pi m\alpha-\nu)g(m)=\varepsilon g(m) \quad (3)$$

by using the ansatz $\psi(ma,\,na)=\mathrm{e}^{\mathrm{i}\nu n}g(m)$, where $x=ma$, $y=na$, and $\varepsilon=-E/V_0$. Solving the eigenvalue problem for specific, i.e., rational values of the magnetic flux $\alpha=\frac{\mathrm{e}a^2B}{hc}$ per elementary plaquette makes the problem periodic. This results in a band spectrum whose gaps form the famous Hofstadter butterfly (Fig. 1). Note that the regime of this spectrum requires finite values of $\alpha$, i.e., magnetic fields $B\sim 1/a^2$, which in the continuum limit $a\to 0$ become ultra-intense. A physical system satisfying (3) can be realized in a simpler way in a 1D optical lattice setup with standard tunneling and an overlaid super-lattice potential with spatial period $1/\alpha$. This statement also holds for the non-Abelian generalizations in the scope of this manuscript. Anyhow, the full 2D realization proposed in [5] allows for a more powerful control opening up completely novel features in such a system.

We consider an atomic gas in a 3D optical lattice and assume that tunneling is completely suppressed in the $z$-direction, so that, effectively, we deal with an array of 2D lattice gases and are able to restrict ourselves to one copy. The atoms occupy two Internal hyperfine states $|g\rangle$, $|e\rangle$, and the optical potential traps them in the state $|g\rangle$ and $|e\rangle$, respectively, in every second column, i.e., for the $y$ coordinate equal to $\ldots,n-1,n+1,\ldots$ ($\ldots,n,n+2,\ldots$. The resulting 2D-lattice has thus the spacing $\lambda/2$ ($\lambda/4$) in the $x$- ($y$-) direction. The tunnel rates in the $x$ direction are due to kinetic energy; they are spatially homogeneous and assumed to be equal for both hyperfine states. The lattice is tilted in the $y$-direction, which introduces an energy shift $\Delta$ between neighboring columns. Tilting can be achieved by accelerating the lattice, or by placing it in a static electric field linearly dependent on $y$.

By this, standard tunneling rates due to kinetic energy are suppressed in the $y$ direction. Instead, tunneling is laser assisted, and driven by two pairs of lasers resonant for Raman transitions between $|g\rangle$ and $|e\rangle$, i.e., $n\leftrightarrow n\pm 1$. This can be achieved because the offset energy for both transitions is different and equals $\pm\Delta$. Detunings of the lasers are chosen in such a way that the effect of tilting is cancelled in the rotating frame of reference. The lasers consist of running waves in the $\pm x$-direction, so that the corresponding tunneling rates acquire local phases $\exp(\pm iqx)$.

In order to realize "artificial" non-Abelian fields in a similar scheme, one has to use atoms with internally degenerate Zeeman sublevels in the hyperfine ground state manifolds, $|g_i\rangle$, and $|e_i\rangle$ with $i=1,\ldots,n$, whose degeneracy is lifted in external magnetic fields. These internal states may be thought of as "colors" of the gauge fields. Promising fermionic candidates with these properties are heavy Alkali atoms, for instance, $^{40}$K atoms in states $F=9/2, M_F=9/2, 7/2,\ldots$, and $F=7/2, M_F=-7/2,-5/2,\ldots$; in particular, they allow for realizing "spin" dependent lattice potentials and spin dependent hopping [10].

Having identified the "colors", one should essentially repeat the scheme of Ref. [5] with two additional modifications: laser assisted tunneling rates along the $y$-axis should depend on the internal state, although not necessarily in the sense of Ref. [10]. They rather should be performed in such a way that for a given link $|g_i\rangle$ to $|e_i\rangle$, tunneling should be described by a non-trivial unitary tunneling matrix $U_y(x)$ being a member of the "color" group ($U(n)$, $SU(n)$, $GL(n)$ etc.). For unitary groups, the tunneling matrix $U_y(x)$ can be represented as $\exp(i\tilde{\alpha}A_y(x))$. Here, $\tilde{\alpha}$ is real, and $A_y(x)$ is a Hermitian matrix from the corresponding gauge algebra, e.g., $\mathfrak{u}(n)$ or $\mathfrak{su}(n)$. Since transitions from $|g_i\rangle$ to $|e_i\rangle$ correspond to different frequencies for each $i$, they are driven by different running wave lasers. Thus, they may attain different phase factors $\exp(\pm iq_ix)$.

In order to create gauge potentials that cannot simply be reduced to two independent Abelian components, tunneling in the $x$-direction should also be Raman laser assisted and should allow for coherent transfer between internal Zeeman states. We will assume for now that this is proceeded by the same unitary tunneling matrix $U_x$ for both hyperfine state manifolds, although more general situations are feasible and basically interesting. To assure a genuine non-Abelian character of the fields, it is necessary that $[U_x, U_y(x)]\neq 0$. We stress that all

FIG. 2: *Optical lattice setup for U(2) gauge fields:* Red and blue open semi-circles (closed semi-circles) denote atoms in states $|g_1\rangle$ and $|g_2\rangle$, respectively ($|e_1\rangle$ and $|e_2\rangle$). Top) Hopping in the $x$-direction is laser assisted and allows for unitary exchange of colors; it is described by the same unitary hopping matrix $U_x$ for both $|g_i\rangle$ and $|e_i\rangle$ states. Hopping along the $y$-direction is also laser assisted and attains "spin dependent" phase factors. Bottom) Trapping potential in $y$-direction. Adjacent sites are set off by an energy $\Delta$ due to the lattice acceleration, or a static inhomogeneous electric field. The lasers $\Omega_{1i}$ are resonant for transitions $|g_{1i}\rangle \leftrightarrow |e_{2i}\rangle$, while $\Omega_{2i}$ are resonant for transitions between $|e_{1i}\rangle \leftrightarrow |g_{2i}\rangle$ due to the offset of the lattice sites. Because of the spatial dependence of $\Omega_{1,2}$ (running waves in $\pm x$ direction) the atoms hopping around the plaquette get the unitary transformation $\hat{U} = U_y^\dagger(m)U_xU_y(m+1)U_x^\dagger$, where $U_y(m) = \exp(2\pi i m\, \mathrm{diag}[\alpha_1, \alpha_2])$, as indicated in upper figure.




elements of our scheme, as shown in Fig. 2, are experimentally accessible. Nevertheless, consistent gauge group realizations demand tunneling matrix amplitudes to be controlled in such a way that they strictly belong to the corresponding unitary groups. which assures the vector potentials to be composed of the Hermitian generators[18]. The scheme of Fig. 2 allows to generalize the Hamiltonian (1) to the case of non-Abelian vector potentials. In fact, we replace the components of $\vec{A}$ by the corresponding matrices from the group algebra. In particular, the illustrated setup generates "artificial" gauge potentials of the form $\vec{A} = (A_x, A_y(m), 0)$, with

$$\vec{A}(m,n) = \frac{\hbar c}{ea}\left(\begin{pmatrix} 0 & \frac{\pi}{2}\mathrm{e}^{\mathrm{i}\phi} \\ \frac{\pi}{2}\mathrm{e}^{-\mathrm{i}\phi} & 0 \end{pmatrix}, \begin{pmatrix} 2\pi m\alpha_1 & 0 \\ 0 & 2\pi m\alpha_2 \end{pmatrix}, 0\right). \quad (4)$$

More precisely, the scheme associates a unitary tunneling operator with every link in analogy with standard lattice gauge theory prescriptions [12]: $U(m-1,n \to m,n) \equiv U_x$, $U(m,n \to m-1,n) \equiv U_x^\dagger$, $U(m,n \to m,n+1) \equiv U_y(m)$, $U(m,n \to m,n-1) \equiv U_y^\dagger(m)$, where $U_x = \exp(+ieaA_x/c\hbar)$ and $U_y = \exp(+ieaA_y(m)/c\hbar)$. The only difference is that $\vec{A}$ acquires an overall factor of $\hbar c/ea$; in effect, though it does not behave well in the continuous limit $a \to 0$, the "magnetic flux" per plaquette, $\alpha_{1,2}$ remains finite. Thus, we are in the same limit of ultra-intense fields $\sim 1/a^2$ as in the "classic" Hofstadter case of Abelian magnetic fields. The ansatz $\psi(ma,\, na) = \mathrm{e}^{\mathrm{i}\nu n}g(m)$, leads to a generalized Harper wave equation

$$\begin{pmatrix} g(m+1) \\ g(m) \end{pmatrix} = B(m)\begin{pmatrix} g(m) \\ g(m-1) \end{pmatrix} \quad (5)$$

with

$$B(m) = \begin{pmatrix} \begin{pmatrix} 0 & \mathrm{e}^{\mathrm{i}\phi}\varepsilon_m(\alpha_2,\,\nu) \\ \mathrm{e}^{-\mathrm{i}\phi}\varepsilon_m(\alpha_1,\,\nu) & 0 \end{pmatrix} & \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix},$$

where $\varepsilon_m(\alpha,\,\nu) = \varepsilon - 2\cos(2\pi m\alpha - \nu)$ is the Harper energy term. Both, $\psi(m,\,n)$ and $g(m)$ are now two-component objects. In the particular case of eq. (4), when two successive transfer matrices $B(m)$ are multiplied by each other, eq. (5) decomposes into a pair of equivalent 2D equations. Nevertheless, to obtain the spectrum, one has to rely on numerical methods. Given each $\alpha_i = p_i/q_i$ rational, the problem is $Q$-periodic (where $Q$ equals the smallest common multiple of $q_1$ and $q_2$). The allowed energies are those for which the product of $Q$ successive matrices $B(m)$ has an eigenvalue of modulus 1. The spectrum shows a band structure, is continuous, and is bounded by two hyperplanes (depicted in grey in Fig. 3). It exhibits a very complex formation of holes of finite measure and various sizes, which we name the Hofstadter "moth". Although a rigorous proof can not be provided, the "moth" reminds of a fractal structure. Obviously, this fractal structure will be very sensitive to any sort of perturbation (finite size of the system, external trapping potential etc.) on very small scales. But, since the holes are true 3D objects with finite volume, the spectrum will be more robust on a larger scale to perturbations than in case of the Hofstadter "butterfly". This may be revealed in comparison of our "moth" with two uncoupled Abelian "butterflies". The latter would form a spectrum of allowed energies consisting of intersecting lamellas of zero width. Here, the perturbations are required to guarantee allowed regions of finite volume. Since $\alpha_1$ and $\alpha_2$ are easily controllable, the spectral structure is directly measurable; a possible scheme would be to load a dilute (weakly interacting, or practically non-interacting) Bose condensate into the lattice and measure its excitations, e.g., by looking at the time evolution of the particle density, as suggested in Ref. [5]. Alternatively, one could load an ultra-cold polarized Fermi gas and measure its Fermi energy as a function of the number of particles.

It is interesting to consider yet another effect that becomes particularly spectacular in the limit of ultra-intense gauge fields, and that can be measured in the proposed system: a non-Abelian Aharonov-Bohm effect, to be considered as an example of non-Abelian interference. In order to realize it, one should prepare, for instance, a weakly interacting Bose

FIG. 3: The Hofstadter "moth" spectrum. Eigenenergies $\varepsilon$ are plotted versus $\alpha_i = p_i/q_i, \in [0, = 0.5]$ $(i = 1, 2)$, where $q_i \leq 41$ and $\alpha_1 \neq \alpha_2$.




condensate in a definite internal state $|\psi_0\rangle$ around a location $P_1$. Then, the BEC (or parts of it) should be split by Raman scattering, and piecewise dragged (using, e.g, laser tweezers) to a meeting point $P_2$ on two distinct paths. These correspond to the unitary transporters $U_1$ and $U_2^\dagger$, respectively. A measurement of the density of atoms at $P_2$ will reveal a non-Abelian interferometer signal, i.e., it will detect the interference term $n_{int} \propto \langle\Psi_0|U_2U_1|\psi_0\rangle$. Choosing, e.g., the rectangular loop, consisting of $L_{x_1}$ steps in the $-x$-direction, $L$ steps in the $y$-direction, $L_{x_1}+L_{x_2}$ steps in $x$-direction, $L$ steps in $-y$ directions, and finally $L_{x_2}$ steps in $-x$-direction, we obtain for the gauge potential of eq. (4): $n_{int} \propto 1$ if $L_{x_1} + L_{x_2}$ is even, and $n_{int} \propto \langle\Psi_0| \exp[\pm 2\pi i\hat{\alpha}L(L_{x_1} + L_{x_2})]|\psi_0\rangle$, if $L_{x_1} + L_{x_2}$ is odd, where $\hat{\alpha} = \mathrm{diag}(\alpha_1, \alpha_2)$. The signal is thus extremely sensitive to $L_{x_1}$ and $L_{x_2}$. If one attempted to measure the phase shifts by introducing obstacles on the $y$ arms of this interferometer, the result would strongly depend on the $x$-coordinate of the obstacles, and the location of $P_1$, $P_2$; a non-Abelian manifestation of the external gauge potential.

Obviously, the properties of the considered system in the limit of ultra-intense fields are quite complex. Though, to get a better intuition concerning the scope of this scheme, it is useful to consider also the "continuum" limit $a \to 0$ with $V_0 \to a^2/m$. Then, the Hamiltonian becomes $\mathcal{H} = (\vec{p} - e/c\vec{A})^2/2m$. A natural question to ask is what kinds of gauge fields of "normal", i.e., $a$-independent strength, can be realized? In other words: what are the possible "artificial" gauge potentials that can be created? In general, phase factors resulting from running wave vectors can be introduced for all tunneling matrices: $\vec{A}(\vec{r}) = \frac{c}{el}(M_1 + [M_2(\frac{x}{l}) + M_3(\frac{y}{l})], N_1 + [N_2(\frac{x}{l}) + N_3(\frac{x}{l})], 0)$, where $M_i, N_i$ are arbitrary (in general non-commuting), dimensionless, and $a$-independent matrices from, e.g., $\mathfrak{u}(2)$, and $l$ is the characteristic length on which $\vec{A}$ varies. Note that in order to realize non-Abelian interferometry, it is sufficient to fulfill $[M_1, N_1] \neq 0$ and set all other matrices to zero. This may be achieved using simpler experimental arrangements as those discussed above, and will be the subject of further investigation. Furthermore, local disorder may be introduced in a controlled way that allows for small fluctuations of the matrices $M_i$. In particular, disorder can be made annealed, i.e. it changes on a time scale comparable with the relevant time scales of the system, and, thus, mimics thermal fluctuations. It can be of significant amplitude, provided it does not drive the assisting lasers system out of resonance. For example, wave vectors of running waves can strongly fluctuate and force the magnetic fluxes to fluctuate correspondingly. Even more complicated spatial dependences, e.g., piecewise linearity, of $\vec{A}$ are feasible by using static electric fields, laser induced potentials, etc. Additional lasers may introduce local, and in general time dependent unitaries. Such transformations would generate arbitrary local temporal components of the gauge potential, $A_0(x, y)$. Although for Yang-Mills fields in 2+1D, this component may be gauged out adapting the Weyl, or strict temporal gauge [19], the corresponding gauge transformations may introduce more complex spatial and temporal forms of the remaining two components of $\vec{A}$.

Finally, let us discuss, in which sense the proposed 2D scheme might be useful to study lattice gauge theories (LGT) in 2+1D. There, one uses the framework of Euclidean field theory, and various methods of statistical physics, such as Monte Carlo methods, to sum over all configurations of gauge and matter fields at finite or infinite temperatures. Thus, gauge fields are dynamical variables in LGT, but are obviously not in the proposed scheme. Moreover, our scheme is realized in real rather than in imaginary time, thus, only a limited set of configurations of gauge fields can be generated. Nevertheless, the big advantage of our proposal is that given a gauge field configuration, the dynamics of matter fields in real time are given for free. When experimentally realized, the system may be used as a quantum simulator of matter fields in a given external gauge field. By generating other configurations, we may try to "mimic" the Monte Carlo sampling of LGT. Averaging over both, annealed disorder and quantum fluctuations should approximate the statistical average in LGT. However, this inevitably requires that generated configurations represent the characteristic or statistically relevant ones of corresponding LGT phases.

Apparently, with the limited gauge field configurations accessible in the proposed scheme, not too much can be done. Anyhow, at least some configurations may be generated that share characteristics of LGT phases, e.g., an area law fulfilled by Wilson loops in the confinement sector. In fact, the $SU(2)$ gauge potential of eq. (4) with fluctuating, but anti-correlated fluxes $\alpha_1 = -\alpha_2$ yields an area law for Wilson loops in the $xy$-plane, provided the probability distribution of the fluxes is Lorentzian. However, it would be desirable to create configurations that exhibit other characteristics of the confinement phase such as appropriate distributions of center vortices, Abelian magnetic monopoles, instantons, merons, calorons etc. [11, 20].

We expect that this program will lead to many fascinating results. It should be stressed that Yang-Mills theories in 2+1D

are in the center of interest in high energy physics, as they describe the high temperature behavior of 4D models [21]. Recently, there has been progress in understanding these theories [19, 22] in the continuum limit, and in the pure gauge sector: the gauge-invariant degrees of freedom have been identified and the Hamiltonian has been constructed. The ground state wave function is, to a very good approximation, a Gaussian function of the currents, and quantities as the string tension are known exactly. Our lattice models may shed new light on these recent discoveries.

We acknowledge fruitful and inspiring discussions with H.-J. Diehl, O. Lechtenfeld, H. Frahm, F. Haake, C. Sämann, and H. Schulz. This work was supported by the Deutsche Forschungsgemeinschaft (SPP1116, SFB407, 436POL), the ESF PESC QUDEDIS, and the Spanish MEC grant HA2002-0036.

[1] N. K. Wilkin and J. M. F. Gunn, Phys. Rev. Lett. **84**, 6 (2000). B. Paredes, P. Fedichev, J. I. Cirac, and P. Zoller , Phys. Rev. Lett. **87**, 010402 (2001); B. Paredes, P. Zoller, and J. I. Cirac, Solid State Comm. **127**, 155 (2003).

[2] P. Rosenbusch *et al.*, Phys. Rev. Lett. **88**, 250403 (2002); V. Bretin, S. Stock, Y. Seurin, and J. Dalibard, Phys. Rev. Lett. **92**, 050403 (2004); JILA rotating trap

[3] M. A. Baranov, *et al.*, in *Condensation and Coherence in Condensed Matter*, Proc. Nobel Jubilee Symposium, Göteborg 2001, Eds. T. Claeson and P. Delsing, Physica Scripta, v. T102, 74 (2002).

[4] M. A. Baranov, K. Osterloh, and M. Lewenstein, cond-mat/0404329, Phys. Rev. Lett. **94** in print (2005).

[5] D. Jaksch and P. Zoller, New J. Phys. **5**, Art. 56 (2003).

[6] E. J. Mueller, cond-mat/0404306.

[7] A. S. Sorensen, E. Demler, M. D. Lukin, cond-mat/0405079

[8] G. Juzeliunas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).

[9] D. R. Hofstadter, Phys. Rev. B**14**, 2239 (1976).

[10] W. V. Liu, F. Wilczek, and P. Zoller, Phys. Rev. A **70**, 033603 (2004)

[11] For $n = 2$ the $U(2)$ lattice gauge theory is equivalent to $SU(2) \times U(1)$ gauge theory; c.f. M. Baig and J. Colet, Phys. Lett. B**176**, 149 (1986).

[12] I. Montvay and G. Münster, *Quantum Fields on a Lattice*, (Cambridge University Press, Cambridge, 1994); H. J. Rothe, *Lattice Gauge Theories* (World Scientific, Singapore, 1998).

[13] N. Craigie (Ed.), *Theory and Detection of Magnetic Monopoles in Gauge Theories*, (World Scientific, Singapore, 1986).

[14] A. Shapere and F. Wilczek, *Goemetric Phases in Physics*, (World Scientific, Singapore, 1989).

[15] P. Öhberg, M. Fleischhauer, private communication.

[16] B. Damski *et al.*, Phys. Rev. Lett. **90**, 110401 (2003).

[17] P. G. Harper, Proc. Phys. Soc. Lond. A**68**, 874 (1955).

[18] In general tunneling matrices could belong to $GL(n)$; such models are of interest themeselves.

[19] For a very good pedagogical review of Yang-Mills systems in 2+1D, see H. Schulz, hep-ph/0008239.

[20] For a recent review of LGT's see M. Engelhardt, hep-lat/0409023; J. Greensite, Prog. Part. Nucl. Phys. **51**, 1 (2003).

[21] S. Deser, R. Jackiw, and S. Templeton, Ann. Phys. **140**, 372 (1982).

[22] D. Karabali, C. Kim, and V. P. Nair, Nucl. Phys. B**524**, 661 (1998).